

BRONICA
CANON CAMERAS
AND BINOCULARS
CONTAX

DELTA INTERNATIONAL
For all your photographic needs...and more!
877-782-9840 deltainternational.com [CLICK HERE](#)



SCIENCE NEWS CATEGORIES
OUR SOLAR SYSTEM
OTHER SOLAR SYSTEMS
STARS
GALAXIES
THE MILKY WAY

COSMOLOGY
SPACE MISSIONS
OBSERVATORIES
LIFE IN THE UNIVERSE
PEOPLE



Try two issues free! Click here

Login Logout
Register
Update Profile

Astronomy Magazine

News
Science
Hobby
Feature Stories
Science
Hobby

Photo Gallery
Astroshops

TheSky Online
For Beginners

Discussion Forums
Astronomy Quiz

Organization Guide
Calendar of Events
Vendor Directory

■ = registered content



SCIENCE NEWS STARS

NOW!

Bursting Beachballs

The explosive collapse of beachball-sized pockets of plasma creates nanosecond pulses in pulsar radio bursts.

by Pamela L. Gay



The Crab Nebula surrounds a pulsar that produces strong radio emissions.
ESO


[Neutron](#) stars are the dense coals of once-massive stars that have given up their atmospheres and collapsed into remnants so dense that protons and electrons are crushed together into neutrons. When neutron stars they can become pulsars — sending periodic blasts of radio emissions toward Earth when their magnetic poles are pointed in our direction. Recent observations using new high-speed detectors have allowed astronomers to discover subpulses within one pulsar's radio bursts. The extremely short duration of these subpulses indicates that they originate from regions no more than two feet across.

With the help of engineers from the National Radio [Astronomy](#) Observatory, a team of astronomers from the New Mexico Institute of Mining and Technology (New Mexico Tech) built specialized electronic equipment to study pulsars at very small timescales. The team used this equipment at the 1000-foot-diameter Arecibo [radio telescope](#) to look for variations in the radio intensity of pulses from the Crab [pulsar](#). Located in the heart of the Crab [Nebula](#), this pulsar spins once every 33 seconds. It is one of three pulsars known to produce extra strong or "giant" pulses of radio [emission](#) interspersed among its normal pulses.

The researchers found that the Crab's "giant" pulses contain subpulses that each last no longer than two nanoseconds. Light travel time mandates that the region producing pulses this short can be no more than about two feet across.

Before this study, three different theories were used to explain the radio pulses of spinning neutron stars. "The small size of these regions is inconsistent with all but one proposed theory for how the radio emission is generated," said team leader Tim Hankins.

He and his colleagues suspects that [density](#) waves in the [plasma](#) at the surface of the [neutron star](#) interact with their own electrical field, becoming progressively denser until they "collapse explosively," creating superstrong bursts of [radio waves](#). The team presents its observations in the March 13 issue of *Nature*.



This diagram shows how a pulsar beams its radiation.
NRAO

Related Stories

- [Stars: High School Students Discover Pulsar](#)
- [Stars: Pulsar Linked to Historic Supernova](#)
- [Stars: Young Supernova Remnant May Hold Pulsar](#)
- [Stars: Pulsar Ages May Be Wrong](#)
- [Stars: Astronomers Hit Pulsar Bulls-Eye](#)

Email-A-Friend

03/16/2003

[Site Map](#) | [Glossary](#) | [Privacy Statement](#) | [Contact Us](#) | [Advertising Info](#) | [About Us](#) | [Employment](#)

Copyright © 1996-2003 Kalmbach Publishing Co.